



Exercise-induced bronchoconstriction alters airway nitric oxide exchange in a pattern distinct from spirometry

Hye-Won Shin, Christina D. Schwindt, Anna S. Aledia, Christine M. Rose-Gottron, Jennifer K. Larson, Robert L. Newcomb, Dan M. Cooper and Steven C. George

Am J Physiol Regulatory Integrative Comp Physiol 291:1741-1748, 2006. First published Jul 13, 2006; doi:10.1152/ajpregu.00178.2006

You might find this additional information useful...

This article cites 57 articles, 35 of which you can access free at:

<http://ajpregu.physiology.org/cgi/content/full/291/6/R1741#BIBL>

Updated information and services including high-resolution figures, can be found at:

<http://ajpregu.physiology.org/cgi/content/full/291/6/R1741>

Additional material and information about *American Journal of Physiology - Regulatory, Integrative and Comparative Physiology* can be found at:

<http://www.the-aps.org/publications/ajpregu>

This information is current as of November 29, 2006 .

Exercise-induced bronchoconstriction alters airway nitric oxide exchange in a pattern distinct from spirometry

Hye-Won Shin,¹ Christina D. Schwindt,^{3,4} Anna S. Aledia,^{1,6} Christine M. Rose-Gottron,⁴ Jennifer K. Larson,^{3,4} Robert L. Newcomb,^{4,5} Dan M. Cooper,^{1,3,4} and Steven C. George^{1,2}

¹Department of Biomedical Engineering, ²Department of Chemical Engineering and Materials Science, ³Department of Pediatrics, ⁴General Clinical Research Center, ⁵Center for Statistical Consulting, and ⁶Department of Medicine, Division of Pulmonary and Critical Care, University of California, Irvine, California

Submitted 14 March 2006; accepted in final form 6 July 2006

Shin, Hye-Won, Christina D. Schwindt, Anna S. Aledia, Christine M. Rose-Gottron, Jennifer K. Larson, Robert L. Newcomb, Dan M. Cooper, and Steven C. George. Exercise-induced bronchoconstriction alters airway nitric oxide exchange in a pattern distinct from spirometry. *Am J Physiol Regul Integr Comp Physiol* 291: R1741–R1748, 2006. First published July 13, 2006; doi:10.1152/ajpregu.00178.2006.—Exhaled nitric oxide (NO) is altered in asthmatic subjects with exercise-induced bronchoconstriction (EIB). However, the physiological interpretation of exhaled NO is limited because of its dependence on exhalation flow and the inability to distinguish completely proximal (large airway) from peripheral (small airway and alveolar) contributions. We estimated flow-independent NO exchange parameters that partition exhaled NO into proximal and peripheral contributions at baseline, postexercise challenge, and postbronchodilator administration in steroid-naïve mild-intermittent asthmatic subjects with EIB (24–43 yr old, $n = 9$) and healthy controls (20–31 yr old, $n = 9$). The mean \pm SD maximum airway wall flux and airway diffusing capacity were elevated and forced expiratory flow, mid-expiratory phase (FEF_{25–75}), forced expiratory volume in 1 s (FEV₁), and FEV₁/forced vital capacity (FVC) were reduced at baseline in subjects with EIB compared with healthy controls, whereas the steady-state alveolar concentration of NO and FVC were not different. Compared with the response of healthy controls, exercise challenge significantly reduced FEV₁ ($-23 \pm 15\%$), FEF_{25–75} ($-37 \pm 18\%$), FVC ($-12 \pm 12\%$), FEV₁/FVC ($-13 \pm 8\%$), and maximum airway wall flux ($-35 \pm 11\%$) relative to baseline in subjects with EIB, whereas bronchodilator administration only increased FEV₁ ($+20 \pm 21\%$), FEF_{25–75} ($+56 \pm 41\%$), and FEV₁/FVC ($+13 \pm 9\%$). We conclude that mild-intermittent steroid-naïve asthmatic subjects with EIB have altered airway NO exchange dynamics at baseline and after exercise challenge but that these changes occur by distinct mechanisms and are not correlated with alterations in spirometry.

asthma; model; inflammation

NITRIC OXIDE (NO) appears in the exhaled breath and performs many functions in the lungs such as smooth muscle relaxation, host defense, inhibition of platelet aggregation, and neurotransmission. Much research effort has focused on utilizing the concentration of NO in the exhaled breath at a constant exhalation flow (C_{ENO}) as a noninvasive marker of inflammation in diseases such as asthma (2, 5). Changes in C_{ENO} during and after exercise have been reported (6, 8, 20, 29, 30, 33, 51), and more recently, alterations in C_{ENO} have been reported to play a role in the

pathogenesis of both exercise-induced (EIB) and thermally-induced bronchoconstriction (9, 21–23, 27, 38, 50).

EIB is thought to be triggered by increased heat and water losses from the airways during exercise (31), leading to airway inflammation and bronchoconstriction. However, even well-controlled steroid-treated asthmatics can experience EIB (4). Furthermore, baseline spirometry (forced expiratory volume in 1 s, FEV₁) cannot predict the presence of EIB (19), and C_{ENO} is elevated at baseline in asthmatics independent of the presence of EIB (7). Thus a complete physiological understanding of alterations in spirometry and C_{ENO} observed in EIB remains unknown.

There is a growing body of evidence that suggests C_{ENO} has both a proximal (i.e., large airway) and peripheral (i.e., small airway and alveolar) contribution (18, 28, 35, 44, 52). This contrasts sharply with other endogenously produced gases, such as carbon dioxide, which are excreted mainly in the alveolar region of the lungs. In healthy adults, $>50\%$ of C_{ENO} arises from the large airways (lobar bronchi and larger) (10, 43). It has been postulated that the elevated C_{ENO} observed in asthma is due to an upregulation of one or more of the nitric oxide (NO) synthase (NOS) isoforms [inducible NOS, endothelial NOS, or neuronal NOS (1, 15, 17, 47, 55, 57)] and a peripheral extension of NO-producing cells in the airways (44). Thus our limited understanding of the role of NO in EIB is due, in part, to the fact that C_{ENO} cannot distinguish proximal and more peripheral contributions.

Our group has previously described a two-compartment (airway and alveolar regions) model of NO exchange (52) and a single-breath technique (53) to estimate flow-independent NO exchange parameters: global maximum airway flux of NO (J'_{awNO}), global airway diffusing capacity (D_{awNO}), airway wall concentration of NO (C_{awNO}), and steady-state alveolar concentration of NO (C_{ANO}) (14). The flow-independent NO exchange parameters can partition C_{ENO} into proximal (J'_{awNO} , D_{awNO} , and C_{awNO}) and peripheral (C_{ANO}) contributions and thus potentially provide insight into the mechanisms of NO exchange in EIB. The goal of the present study was to 1) characterize regional (proximal and peripheral) NO exchange dynamics in steroid-naïve mild asthmatic subjects with EIB at baseline, after exercise challenge, and after bronchodilator administration, and 2) determine whether dynamic changes in NO exchange are correlated with standard indexes of spirometry.

The costs of publication of this article were defrayed in part by the payment of page charges. The article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Address for reprint requests and other correspondence: S. C. George, Dept. of Biomedical Engineering, 3120 Natural Sciences II, Univ. of California, Irvine, CA 92697-2715 (e-mail: scgeorge@uci.edu).



METHODS
Glossary

$A_{I,II}$	Area under the curve in phases I and II of exhaled NO profile (ppb·ml)
C_{ANO}	Mixed or average fractional concentration of NO in gas phase of alveolar region (ppb; a steady-state concentration is achieved for breathhold or exhalation times >10 s)
C_{awNO}	Airway wall concentration of NO (ppb)
$C_{ENO,i}$	Exhaled NO concentration at the mouth at constant exhalation flow i (ppb)
C_{obsNO}	Exhaled NO concentration (ppb) observed by analytical instrument
C_{peakNO}	Maximum or peak exhaled NO concentration (ppb) observed by analytical instrument
D_{awNO}	Diffusing capacity (ml/s) of NO in entire airway tree, expressed as volume of NO per second per fractional concentration of NO in gas phase [$ml\ NO \cdot s^{-1} \cdot (ml\ NO/ml\ gas)^{-1}$] and equivalent to $pl \cdot s^{-1} \cdot ppb^{-1}$
J'_{awNO}	Maximum total volumetric flux of NO from airways ($ppb \cdot ml \cdot s^{-1}$ or pl/s)
$V_{I,II}$	Exhaled volume in phases I and II of exhalation profile (ml)
V_{air}	Volume of airway tree (ml), defined by the subject's ideal body weight (lb) plus age in yr (53), which is very similar to the cumulative volume of airway generations 0–17 based on Weibel (56)
\dot{V}_E	Exhalation flow (ml/s)
W_{50}	Volume (or width) in phases I and II of exhaled NO signal, calculated by taking the volume at which the exhaled concentration is >50% of C_{peakNO}

Subjects

Nine steroid-naive mild-intermittent atopic asthmatic adults with a clinical history of EIB (ages 24–43 yr) and nine healthy adult controls (ages 20–31 yr) participated in this study. Inclusion criteria for the EIB group were a clinical history of mild-intermittent asthma, EIB, and a >10% decrease in FEV₁ following a 10-min exercise challenge (3). Exclusion criteria included a history of smoking, pulmonary diseases other than asthma, cardiovascular or neurological disease, current or previous use of a corticosteroid to manage asthmatic symptoms, or use of a bronchodilator in the 6- to 24-h (depending on the specific bronchodilator) preceding exercise testing. For the healthy adult group, inclusion criteria included no history of heart disease, lung disease, or smoking and normal standard spirometry [FEV₁/forced vital capacity (FVC) > 80% predicted]. The protocol was approved by the Institutional Review Board at the University of California, Irvine, and written informed consent was obtained.

Protocol

Subjects refrained from exercise and food for 72 and 3 h, respectively, before the study. Baseline exhaled NO measurements and spirometry were obtained from each subject. Spirometry included FVC, FEV₁, and forced expiratory flow, midexpiratory phase (FEF_{25–75}) measured in triplicate (V_{max} 229; SensorMedics, Yorba Linda, CA) according to American Thoracic Society (ATS) guidelines (3). Each subject then completed 10 min of exercise challenge (target intensity of 80% of the predicted maximum heart rate at room temperature and ~50% relative humidity) according to ATS guidelines (3), 5 min of recovery, measurement of exhaled NO and spirometry, inhalation of a bronchodilator (3 puffs of Combivent with spacer; Boehringer Ingelheim, Ridgefield, CT), 10 min of rest, and a final measurement of exhaled NO and spirometry. Each puff of Combivent delivers ~18 μg of ipratropium bromide and 103 μg of

albuterol sulfate. Five of the nine subjects with EIB participated in a second control visit in which the exercise period was replaced with a rest period to determine whether spirometry impacted the NO exchange dynamics (24, 45, 49).

Exhaled NO Measurement

A chemiluminescence NO analyzer (Sievers, Boulder, CO) and pneumotachometer (Hans Rudolph, Kansas City, MO) were used to record NO, pressure, and flow for five repetitions in each subject of a 20-s preexpiratory breathhold followed by a decreasing exhalation flow maneuver (41, 53). The profiles were characterized by the peak concentration in phases I and II, C_{peakNO} , the volume (or width) of phases I and II, W_{50} , and the total mass of NO, $A_{I,II}$ (Fig. 1). In addition, with a two-compartment model (14, 52), the profiles were used to determine the flow-independent NO exchange parameters (J'_{awNO} , D_{awNO} , C_{ANO} , and C_{awNO}) as previously defined (14). The flow-independent parameters were then used to predict C^*E_{NO} (asterisk denotes calculated value) at a constant exhalation flow (\dot{V}_E) of 50 ml/s by using the relatively simple expression (53):

$$C^*E_{NO,\dot{V}} = C_{awNO} + (C_{ANO} - C_{awNO}) \cdot \exp(-D_{awNO}/\dot{V}_E) \quad (1)$$

Statistics

Untransformed data were analyzed using repeated-measures analysis of variance to detect differences between groups and within subjects over time. Log transformation did not improve the normality of the data distribution. In addition, contrast analyses were performed to assess differences between baseline and postexercise scores, between baseline and postbronchodilator scores, and between postexercise and postbronchodilator scores for each of the dependent variables. Paired comparison t -tests were employed to evaluate the differences between subjects' baseline scores and their scores at each of the other two time points over time. Statistical significance was considered at $P < 0.05$.

RESULTS

The physical characteristics of the subjects, such as age, height, weight, and ideal body weight, were not different




Fig. 1. Model-independent parameters characteristic of the exhalation profile in phases I and II are defined schematically. C_{obsNO} is the exhaled NO concentration observed experimentally from the analytical instrument (ppb, parts per billion); C_{peakNO} is the maximum concentration of NO in phases I and II; W_{50} is the width of the phase I and II peak calculated by taking the volume at which the exhaled concentration is >50% of C_{peakNO} ; $V_{I,II}$ is the volume of phases I and II; and $A_{I,II}$ is the total mass of NO (area under the curve, shown as a shaded region) in phases I and II. The distinction between phases I and II and phase III is the point of zero slope (minimum point) in the exhalation profile as previously described (53).

between subjects with healthy controls and EIB (Table 1). All subjects were able to complete the 10 min of targeted intensity exercise without complication.

FVC, FEV₁, FEF_{25–75}, and FEV₁/FVC at baseline, postexercise challenge, and postbronchodilator administration are presented in Table 2. At baseline, subjects with EIB had lower FEV₁ (%predicted), FEF_{25–75}, and FEV₁/FVC compared with the healthy controls. FEV₁ (liters) and FVC (liters, %predicted) were not significantly different between healthy controls and subjects with EIB. Postexercise, all four indexes of lung function (FVC, FEV₁, FEF_{25–75}, and FEV₁/FVC) were reduced in the subjects with EIB compared with the response of the healthy adults. Postbronchodilator, FEV₁, FEF_{25–75} and FEV₁/FVC were elevated relative to baseline compared with the response of the healthy controls.

All the subjects were able to complete the single breath maneuver with a 20-s preexpiratory breathhold. Composite NO exhalation profiles for subjects with EIB (baseline, postexercise challenge, and postbronchodilator administration) and healthy controls (only presented baseline) were generated by taking the mean exhaled concentration at equivalent exhaled volume intervals of all subjects in a given group and condition (e.g., EIB postexercise) and are presented in Fig. 2. The exercise challenge and bronchodilator administration did not impact the exhalation NO profile for healthy controls. In contrast, subjects with EIB had an increased concentration of NO in all phases of the exhalation profile. For subjects with EIB, significantly less NO was exhaled postexercise challenge, which can be seen by the reduced peak height in phases I and II, smaller area under the curve (A_{I,II}), and lower NO concentrations in phase III of the exhalation profile. In addition, bronchodilator administration increased the NO in all phases of the exhalation profile.

Mean (SD) C_{peakNO} values for subjects with EIB at baseline, postexercise challenge, and postbronchodilator administration were 136 (104), 94 (53), and 112 (65) parts per billion (ppb), respectively. Mean (SD) A_{I,II} values for subjects with EIB at baseline, postexercise challenge, and postbronchodilator administration were 32,600 (22,400), 24,300 (15,500), and 31,900 (19,600) ppb·ml, respectively. Both C_{peakNO} and A_{I,II} values are significantly smaller postexercise challenge compared with the baseline (*P* < 0.05). Mean (SD) W₅₀ for subjects with EIB at baseline was 206 (52) and was not impacted by the exercise challenge or bronchodilator. Mean (SD) C_{peakNO}, W₅₀, and A_{I,II} values for healthy controls at baseline were 72 (32) ppb, 190 (48) ppb, and 12,822 (6,571) ppb/ml respectively, and were not impacted by the exercise challenge or bronchodilator (see Fig. 2).

The experimentally observed changes in exhaled NO concentration in EIB are reflected in changes in the flow-independent NO exchange parameters. *J'*aw_{NO} and Daw_{NO} were statistically elevated at baseline in subjects with EIB, whereas C_{ANO} and Caw_{NO} were not different compared with healthy controls (Fig. 3). For healthy controls, none of the flow-independent NO exchange parameters were significantly different after the exercise challenge or bronchodilator administration compared with baseline (Fig. 4, A, C, E, and G). *J'*aw_{NO} was significantly decreased (mean change of –35%) after the exercise challenge relative to baseline in all nine subjects with EIB (Fig. 4B). Administration of the bronchodilator significantly increased *J'*aw_{NO} relative to postexercise, but the difference relative to baseline was not statistically significant compared with healthy controls. Daw_{NO}, C_{ANO}, and Caw_{NO} were not statistically altered after exercise or bronchodilator administration (Fig. 4, D, F, and H). For the five subjects who participated in the second visit, the mean (SD) changes in

Table 1. Physical characteristics of subjects

Subject	Sex	Age, yr	Height, m	Weight, kg	BMI, kg/m ²	IBW, kg	V _{air} , ml	Therapies
<i>Healthy adults</i>								
1	M	24	1.83	79.4	23.7	76	191	
2	M	27	1.65	75.3	27.7	64	168	
3	M	27	1.80	83.0	25.6	73	189	
4	F	23	1.57	54.4	22.1	54	143	
5	F	31	1.73	56.2	18.8	64	172	
6	F	22	1.73	81.2	27.1	64	163	
7	F	28	1.68	65.3	23.1	61	162	
8	F	26	1.60	50.8	19.8	56	150	
9	F	20	1.63	63.5	23.9	58	148	
Mean		25.3	1.69	67.7	23.5	63.4	165	
SD		3.39	0.09	12.4	3.02	7.26	17.0	
<i>Adults with EIB</i>								
1	F	29	1.63	62.6	23.6	58.1	157	Albuterol, salmeterol
2	M	36	1.78	103	32.5	71.7	194	Albuterol
3	F	26	1.60	70.3	27.5	55.8	149	Albuterol
4	F	43	1.60	67.6	26.4	56.7	168	Albuterol
5	M	26	1.88	89.8	25.4	78.9	200	Albuterol
6	M	24	1.75	71.7	23.4	69.9	178	Albuterol
7	M	25	1.83	77.6	23.2	75.8	192	Albuterol
8	F	31	1.75	69.9	22.8	64.4	173	Albuterol
9	M	27	1.78	92.5	29.2	71.7	185	Albuterol
Mean		29.7	1.73	78.3	26.0	67.0	177	
SD		6.20	0.10	13.6	3.28	8.58	17.3	

EIB, exercise-induced bronchoconstriction (asthma); M, male; F, female; BMI, body mass index; IBW, ideal body weight; V_{air}, volume of the airway compartment estimated in ml as the sum of the subject's ideal body weight (lb.) plus age (yr.) (53).

Table 2. Spirometry at baseline, postexercise challenge, and postbronchodilator

Subject	FVC, liters			FVC, % predicted			FEV ₁ , liters			FEV ₁ , % predicted			FEF ₂₅₋₇₅ , liters			FEV ₁ /FVC		
	Base, liters	EX, %Δ	BD, %Δ	Base, %	EX, %Δ	BD, %Δ	Base, liters	EX, %Δ	BD, %Δ	Base, %	EX, %Δ	BD, %Δ	Base, liters	EX, %Δ	BD, %Δ	Base	EX, %Δ	BD, %Δ
<i>Healthy adults</i>																		
1	5.5	0.4	3.4	97	0.0	3.1	4.7	-0.8	2.5	96	-1.0	3.1	4.9	1.4	10.9	87	-1.1	-1.1
2	4.5	-0.2	0.0	105	-1.0	0.0	3.8	-1.3	-0.5	103	-1.9	-1.0	3.9	-4.3	-7.1	83	-1.2	2.4
3	5.4	0.4	1.7	97	0.0	2.1	4.4	-0.5	4.6	94	0.0	4.3	4.3	-2.6	10.9	82	-1.2	1.2
4	3.3	3.3	5.1	98	3.1	5.1	3.0	3.3	7.0	102	2.9	6.9	4.4	-2.5	13.5	90	0.0	3.3
5	4.4	0.7	0.7	111	0.9	0.0	3.8	1.6	4.5	113	1.8	4.4	4.3	-0.2	8.9	86	1.2	3.5
6	4.9	-1.0	-1.0	119	-0.8	-0.8	4.0	-3.7	1.0	115	-3.5	0.9	4.1	-11.1	0.7	82	-2.4	2.4
7	4.0	4.0	4.3	105	3.8	3.8	3.4	2.1	0.9	106	1.9	0.9	4.1	-10.0	-13.1	87	-2.3	-4.6
8	3.1	4.6	4.6	88	4.5	4.5	2.6	1.2	10.5	85	2.4	10.6	2.5	-5.1	30.0	84	-2.4	6.0
9	3.7	1.1	0.8	102	1.0	1.0	3.2	-8.8	-8.2	98	-9.2	-8.2	3.5	-28.2	-22.1	89	-12.4	-10.1
Mean	4.3	1.5	2.2	102	1.3	2.1	3.6	-0.8	2.5	101	-0.7	2.4	4.0	-7.0	3.6	86	-2.4	0.3
SD	0.9	2.0	2.2	9.0	2.0	2.2	0.7	3.7	5.3	9.4	3.8	5.3	0.7	15.8	3.0	3.9	3.9	4.9
<i>Adults with EIB</i>																		
1	3.7	-0.8	12.7	104	-1.0	12.5	2.4	-11.9	38.7	77	-11.7	39.0	1.4	-19.6	106	64	-12.5	21.9
2	3.4	-19.9	29.9	67	-19.4	29.9	1.9	-27.7	60.1	45	-28.9	60.0	1.0	-33.7	95.9	55	-7.3	23.6
3	3.1	-37.5	-2.9	90	-37.8	-2.2	2.2	-56.7	0.9	73	-56.2	1.4	1.5	-74.3	8.1	70	-30.0	4.3
4	2.1	-11.1	17.9	64	-10.9	18.8	1.4	-20.0	33.3	50	-20.0	32.0	0.8	-34.2	76.3	65	-9.2	13.8
5	5.4	-16.4	-1.3	85	-16.5	-1.2	3.7	-31.0	18.9	70	-31.4	18.6	2.5	-51.8	75.1	69	-20.3	18.8
6	5.1	-2.1	3.3	114	-2.6	3.5	3.3	-18.2	12.9	93	-18.3	12.9	2.0	-36.5	28.1	63	-15.9	9.5
7	5.6	-3.9	2.5	98	-4.1	2.0	4.1	-10.0	23.8	83	-10.8	22.9	3.1	-14.8	91.6	73	-6.8	20.5
8	3.8	-1.3	-0.5	94	-1.1	0.0	3.0	-11.7	2.3	89	-11.2	2.2	2.8	-39.0	19.1	80	-11.3	2.5
9	5.5	-12.7	-9.1	103	-12.6	-8.7	4.5	-17.1	-6.7	101	-17.8	-6.9	4.6	-27.9	2.0	82	-6.1	2.4
Mean	4.2	-11.8†	5.8	91.0	-11.8†	6.1	2.9	-22.7†	20.5†	75.7*	-22.9†	20.2†	2.2*	-36.9†	55.8†	69*	-13.3†	13.1†
SD	1.3	11.9	12.1	16.7	11.9	12.1	1.1	14.6	21.2	18.8	14.5	21.1	1.2	17.7	41.0	8.5	7.8	8.6

FVC, forced vital capacity; FEV₁, forced expiratory volume in 1 s; FEF₂₅₋₇₅, forced expiratory flow, midexpiratory phase; EX, postexercise challenge; BD, postbronchodilator administration; %Δ, percent change from baseline. *Statistically different from healthy controls at baseline ($P < 0.05$). †Difference between scores relative to baseline is statistically different from healthy controls ($P < 0.05$).

FEV₁ and $J'aw_{NO}$ after the 10-min rest period were -1.4 (1.3)% and -3.5 (18)% and did not represent significant changes from baseline.

Our group previously demonstrated that experimental values of C^*E_{NO} were not different from model-predicted (Eq. 1) C^*E_{NO} for healthy adults (41, 42) and steroid-naive asthmatics (40) at an exhalation flow of 50 ml/s. Thus C^*E_{NO} for subjects with EIB is presented at baseline in Fig. 3B and as a function of time in Fig. 4, I (healthy adults) and J (EIB). Relative to healthy controls, the trend for the changes in C^*E_{NO} in the subjects with EIB was the same as that for $J'aw_{NO}$. For healthy controls, C^*E_{NO} was not

impacted by the exercise challenge or bronchodilator. None of the observed changes in $J'aw_{NO}$ and C^*E_{NO} postexercise and postbronchodilator administration had any significant correlation with changes in spirometric indexes (FVC, FEV₁, FEF₂₅₋₇₅, and FEV₁/FVC) postexercise and postbronchodilator administration for healthy control or EIB subjects.

DISCUSSION

This is the first study to examine the dynamic relationship in EIB between spirometric indexes and proximal and peripheral NO exchange. We found that elevated baseline exhaled NO concentration in subjects with EIB was due primarily to an increase in the airway diffusing capacity of NO, resulting in a larger maximum flux ($J'aw_{NO} = Daw_{NO} \times Caw_{NO}$) of NO from the airway wall. EIB caused a significant decrease in $J'aw_{NO}$ without significant changes in Daw_{NO} or Caw_{NO} . Bronchodilation following EIB returned $J'aw_{NO}$ to near baseline, but this response did not differ significantly from that of healthy controls. In addition, changes in spirometric indexes did not correlate with the airway NO parameters altered in EIB: $J'aw_{NO}$ and Daw_{NO} . We conclude that 1) steroid-naive subjects with EIB have no alterations in alveolar NO but have an elevated baseline airway wall diffusing capacity and maximum airway wall flux of NO; 2) EIB acutely reduces the maximum airway wall flux of NO by a mechanism distinct from the altered baseline NO exchange dynamics; and 3) changes in airway caliber postexercise and postbronchodilator that impact spirometry (FVC, FEV₁, FEF₂₅₋₇₅, and FEV₁/FVC) do not correlate with changes in the NO exchange parameters.




Fig. 2. Composite experimental NO exhalation profiles are presented for the 20-s breathhold followed by a decreasing flow rate maneuver for subjects with asthma [exercise-induced bronchoconstriction (EIB), $n = 9$] at baseline, postexercise (Post-EX) challenge, and postbronchodilator (Post-BD) administration and for healthy controls [healthy adults (HA), $n = 9$] at baseline.




Fig. 3. Mean values of global maximum airway flux of NO (J'_{awNO}), global airway diffusing capacity (D_{awNO}), steady-state alveolar concentration of NO (C_{ANO}), and airway wall concentration of NO (C_{awNO}) with individual data points (A) and mean experimental values of exhaled NO concentration at the mouth at constant exhalation flow (C^*E_{NO}) determined at an exhalation flow of 50 ml/s with individual data points at baseline (B) are presented in subjects with EIB and HA. *Statistically different from healthy controls at baseline.

Before the current study was performed, the relationship between exercise and exhaled NO had been investigated primarily using either exhaled concentration or the product of exhaled concentration and flow (elimination rate). In healthy subjects, exercise causes either no change or a small decrease in the exhaled concentration postexercise, with a large increase in the elimination of NO during exercise due to the increase in ventilation rate (while concentration stays relatively constant) (6, 8, 9, 20, 29, 30, 33, 34, 46, 51). These observed changes last for only a short time (<5 min) postexercise, until ventilation rate returns to baseline. In contrast, exhaled NO concentration in asthma has been reported to be significantly reduced shortly after exercise (9, 48, 50), and the degree of EIB is significantly associated with atopy and baseline exhaled NO (37). These observations are consistent with our current findings in asthmatic and healthy subjects. A major difference between asthmatic and healthy subjects is the observed changes in spirometry following exercise and bronchodilator administration, which reflect changes in the caliber of the airways. Thus region-specific analysis of NO exchange can potentially provide mechanistic insight into the altered NO exchange dynamics between healthy and asthmatic subjects.

There has been only one other study examining region-specific alterations in NO dynamics following exercise; however, this study (39) focused on healthy subjects and a more intense exercise challenge (20 min at 90% of the maximum

heart rate). In this case, D_{awNO} acutely (3 min postexercise) increased, both C_{awNO} and J'_{awNO} decreased, and there was no change in spirometry or C_{ANO} . The decrease in C_{awNO} was attributed to enhanced losses of NO in the exhaled air during the period of exercise, or in other words, a washout of tissue NO stores. These changes were short-lived, and there was no difference from baseline at 30 min postexercise.

At baseline, our data demonstrate that C_{ANO} in subjects with EIB is not different from that in healthy controls. In contrast, J'_{awNO} is approximately fivefold higher in subjects with EIB. Most of this increase can be attributed to an increase (~3-fold) in D_{awNO} and a modest increase in C_{awNO} . Both J'_{awNO} and D_{awNO} are proportional to the airway surface area emitting NO (39, 41). Thus the increase in J'_{awNO} and D_{awNO} at baseline may be due to the peripheral extension of nonadrenergic, noncholinergic nerves from the large airways into the smaller airways (44) or the enhanced expression of inducible NOS in the airway epithelium (16), both of which may increase the airway surface area emitting NO. This pattern in the flow-independent NO parameters (i.e., significant elevation of D_{awNO} and J'_{awNO} with no change in C_{awNO} and C_{ANO}) at baseline is the same as that previously reported in steroid-naive asthmatic subjects (40, 44).

After an exercise challenge, all subjects with EIB demonstrated a marked decrease in only J'_{awNO} and no significant change in C_{ANO} , D_{awNO} , or C_{awNO} . This finding is consistent with a




Fig. 4. Percent changes of J'_{awNO} (A and B), D_{awNO} (C and D), C_{awNO} (E and F), C_{ANO} (G and H), and C^*_{ENO} (I and J; Eq. 1) at postexercise challenge and at postbronchodilator administration relative to baseline are shown in 9 healthy adult subjects (A, C, E, G, and I) and in 9 subjects with EIB (B, D, F, H, and J). Gray shading indicates the window of time for either the exercise challenge or the delivery of bronchodilator; horizontal solid bars indicate the mean value at each time point. #Difference between scores relative to baseline is statistically different from that of healthy controls ($P < 0.05$). ‡Difference between scores relative to postexercise challenge is statistically different from that of healthy controls ($P < 0.05$).

decrease in the airway (proximal) contribution of exhaled NO and no change in the alveolar (peripheral) contribution. Although most subjects demonstrated a mild decrease in D_{awNO} and C_{awNO} (Fig. 3, B and D), some demonstrated an increase. Thus the total airway flux of NO is decreased due to a combination of changes in D_{awNO} and C_{awNO} , and a washout of NO in the airway wall tissue cannot fully explain the observations. The observation that D_{awNO} is altered at baseline but not in response to exercise leads us to conclude that alterations in NO exchange dynamics at baseline are due to mechanisms different from those that alter NO exchange following exercise.

As mentioned earlier, bronchoconstriction following exercise and bronchodilation following bronchodilator administration are major differences between the response of asthmatic and healthy subjects. These changes in airway caliber, which reflect the dynamic changes in FVC, FEV₁, FEF_{25–75}, and FEV₁/FVC, can impact exhaled NO in several ways. First, changes in the caliber of the airways impact the surface area of the airways (26). However, the observation that only J'_{awNO} is altered postexercise and postbronchodilator, when both J'_{awNO} and D_{awNO} are proportional to surface area, suggests that determinants other than surface area are involved and that physical determinants of spirometry are decoupled with NO exchange dynamics.

Second, metabolic determinants may modulate exhaled NO and EIB, such as the production rate from NOS isoforms (9, 48), breakdown of S-nitrosothiols (11, 12, 36), the presence of eosinophils at sites of inflammation (25, 58), or altered mucus production and hydration (39). The later may impact not only the thickness of the diffusion barrier for NO but also the molecular diffusivity (the relative ease with which a molecule can diffuse through a medium), as previously described (39). Bronchoconstriction may occur preferentially at sites of greater inflammation (and thus higher NO production and release), leading to reduced ventilation or air trapping in these areas and a reduction in exhaled NO. This possibility is consistent with the observed mild decrease in FVC (and thus air trapping) following the exercise challenge.

Third, changes in airway caliber that lead to changes in spirometry are complex and only partially understood. Decreases in all of the spirometric indexes can reflect an increased resistance to expiratory air flow due to changes in either the proximal or peripheral airways without providing region-specific information regarding airflow limitation (13, 32). Most recently, it was demonstrated that bronchoconstriction and air trapping in asthma are the result of heterogeneous changes (contraction and dilation) in the airway caliber throughout the airway tree (26, 54). In contrast, exhaled NO may be predominantly from the larger airways (10, 43), thus weakening the relationship with spirometry.

Our two-compartment model makes a simplifying assumption that V_{air} does not change in response to EIB and the bronchodilator. Evidence supporting this assumption is the finding that W_{50} did not change in response to EIB or the bronchodilator. Nonetheless, we have previously reported (53) that only the estimation of D_{awNO} is significantly impacted (an overestimation in V_{air} results in an overestimation in D_{awNO}) by the estimate of V_{air} . Thus this interaction could potentially increase the estimated value of D_{awNO} following EIB. The mean value that we reported for D_{awNO} following EIB was slightly lower compared with baseline, but this change was not significant. However, if V_{air} was overestimated at this time

point, D_{awNO} would be overestimated and could potentially be further from the baseline.

In conclusion, we have quantified several flow-independent parameters characteristic of NO exchange in response to EIB and reported their dynamic relationship with spirometric indexes. The source of elevated exhaled NO at baseline in subjects with EIB is an elevated airway diffusing capacity that increases the airway flux of NO. There is a significant decrease in the airway wall flux of NO after EIB, and bronchodilation returns the airway wall flux to baseline without altering airway diffusing capacity and airway wall concentration. These observations do not correlate with changes in airway caliber. Thus structural changes in the airways that alter spirometric indexes (e.g., FEV₁) cannot completely account for changes in airway NO exchange dynamics. We conclude that elevated exhaled NO at baseline in subjects with EIB and reduced exhaled NO in acute EIB occur by distinct mechanisms, do not correlate with changes in spirometry, and thus reflect both anatomic and metabolic determinants of NO exchange.

ACKNOWLEDGMENTS

We thank the General Clinical Research Center at University of California, Irvine.

GRANTS

This work was supported by National Institutes of Health Grants HL070645, HD23969, ES014923, and RR00827.

REFERENCES

- Adachi H, Nguyen PH, Belardinelli R, Hunter D, Jung T, and Wasserman K. Nitric oxide production during exercise in chronic heart failure. *Am Heart J* 134: 196–202, 1997.
- Alving K, Weitzberg E, and Lundberg JM. Increased amount of nitric oxide in exhaled air of asthmatics. *Eur Respir J* 6: 1368–1370, 1993.
- American Thoracic Society. Guidelines for methacholine and exercise challenge testing-1999. *Am J Respir Crit Care Med* 161: 309–329, 2000.
- Anderson SD. Exercise-induced asthma in children: a marker of airway inflammation. *Med J Aust* 177, Suppl: S61–63, 2002.
- Barnes PJ and Kharitonov SA. Exhaled nitric oxide: a new lung function test. *Thorax* 51: 233–237, 1996.
- Bauer JA, Wald JA, Doran S, and Soda D. Endogenous nitric oxide in expired air: effects of acute exercise in humans. *Life Sci* 55: 1903–1909, 1994.
- Carraro S, Corradi M, Zanconato S, Alinovi R, Pasquale MF, Zaccchello F, and Baraldi E. Exhaled breath condensate cysteinyl leukotrienes are increased in children with exercise-induced bronchoconstriction. *J Allergy Clin Immunol* 115: 764–770, 2005.
- Chirpaz-Oddou MF, Favre-Juvin A, Flore P, Eterradossi J, Delaire M, Grimbert F, and Therminarias A. Nitric oxide response in exhaled air during an incremental exhaustive exercise. *J Appl Physiol* 82: 1311–1318, 1997.
- De Gouw HW, Marshall-Partridge SJ, Van Der Veen H, Van Den Aardweg JG, Hiemstra PS, and Sterk PJ. Role of nitric oxide in the airway response to exercise in healthy and asthmatic subjects. *J Appl Physiol* 90: 586–592, 2001.
- DuBois AB, Kelley PM, Douglas JS, and Mohsenin V. Nitric oxide production and absorption in trachea, bronchi, bronchioles, and respiratory bronchioles of humans. *J Appl Physiol* 86: 159–167, 1999.
- Fang K, Johns R, Macdonald T, Kinter M, and Gaston B. S-nitroso-glutathione breakdown prevents airway smooth muscle relaxation in the guinea pig. *Am J Physiol Lung Cell Mol Physiol* 279: L716–L721, 2000.
- Gaston B, Sears S, Woods J, Hunt J, Ponaman M, McMahon T, and Stamler JS. Bronchodilator S-nitrosothiol deficiency in asthmatic respiratory failure. *Lancet* 351: 1317–1319, 1998.
- Gelb AF, Williams AJ, and Zamel N. Spirometry. FEV₁ vs FEF_{25–75} percent. *Chest* 84: 473–474, 1983.
- George SC, Hogman M, Permutt S, and Silkoff PE. Modeling pulmonary nitric oxide exchange. *J Appl Physiol* 96: 831–839, 2004.

15. **Grasemann H, Yandava CN, Storm van's Gravesande K, Deykin A, Pillari A, Ma J, Sonna LA, Lilly C, Stampfer MJ, Israel E, Silverman EK, and Drazen JM.** A neuronal NO synthase (NOS1) gene polymorphism is associated with asthma. *Biochem Biophys Res Commun* 272: 391–394, 2000.
16. **Hamid Q, Springall DR, and Riveros-Moreno V.** Induction of nitric oxide synthase in asthma. *Lancet* 342: 1510–1513, 1993.
17. **Hansel TT, Kharitonov SA, Donnelly LE, Erin EM, Currie MG, Moore WM, Manning PT, Recker DP, and Barnes PJ.** A selective inhibitor of inducible nitric oxide synthase inhibits exhaled breath nitric oxide in healthy volunteers and asthmatics. *FASEB J* 17: 1298–1300, 2003.
18. **Hogman M, Drca N, Ehrstedt C, and Merilainen P.** Exhaled nitric oxide partitioned into alveolar, lower airways and nasal contributions. *Respir Med* 94: 985–991, 2000.
19. **Holzer K, Anderson SD, and Douglass J.** Exercise in elite summer athletes: challenges for diagnosis. *J Allergy Clin Immunol* 110: 374–380, 2002.
20. **Iwamoto J, Pendergast DR, Suzuki H, and Krasney JA.** Effect of graded exercise on nitric oxide in expired air in humans. *Respir Physiol* 97: 333–345, 1994.
21. **Kanazawa H, Asai K, Hirata K, and Yoshikawa J.** Vascular involvement in exercise-induced airway narrowing in patients with bronchial asthma. *Chest* 122: 166–170, 2002.
22. **Kanazawa H, Hirata K, and Yoshikawa J.** Involvement of vascular endothelial growth factor in exercise induced bronchoconstriction in asthmatic patients. *Thorax* 57: 885–888, 2002.
23. **Kanazawa H, Hirata K, and Yoshikawa J.** Role of endogenous nitric oxide in exercise-induced airway narrowing in patients with bronchial asthma. *J Allergy Clin Immunol* 106: 1081–1087, 2000.
24. **Kissoon N, Duckworth LJ, Blake KV, Murphy SP, and Lima JJ.** Effect of β_2 -agonist treatment and spirometry on exhaled nitric oxide in healthy children and children with asthma. *Pediatr Pulmonol* 34: 203–208, 2002.
25. **Kivity S, Argaman A, Onn A, Shwartz Y, Man A, Greif J, and Fireman E.** Eosinophil influx into the airways in patients with exercise-induced asthma. *Respir Med* 94: 1200–1205, 2000.
26. **Kotaru C, Coreno A, Skowronski M, Muswick G, Gilkeson RC, and McFadden ER Jr.** Morphometric changes after thermal and methacholine bronchoprovocations. *J Appl Physiol* 98: 1028–1036, 2005.
27. **Kotaru C, Skowronski M, Coreno A, and McFadden ER Jr.** Inhibition of nitric oxide synthesis attenuates thermally induced asthma. *J Appl Physiol* 91: 703–708, 2001.
28. **Lehtimäki L, Kankaanranta H, Saarelainen S, Hahtola P, Jarvenpää R, Koivula T, Turjanmaa V, and Moilanen E.** Extended exhaled NO measurement differentiates between alveolar and bronchial inflammation. *Am J Respir Crit Care Med* 163: 1557–1561, 2001.
29. **Maroun MJ, Mehta S, Turcotte R, Cosio MG, and Hussain SN.** Effects of physical conditioning on endogenous nitric oxide output during exercise. *J Appl Physiol* 79: 1219–1225, 1995.
30. **Matsumoto A, Hirata Y, Momomura S, Fujita H, Yao A, Sata M, and Serizawa T.** Increased nitric oxide production during exercise. *Lancet* 343: 849–850, 1994.
31. **McFadden ER Jr and Gilbert IA.** Exercise-induced asthma. *N Engl J Med* 330: 1362–1367, 1994.
32. **McFadden ER Jr, Ingram RH Jr, Haynes RL, and Wellman JJ.** Predominant site of flow limitation and mechanisms of postexercise asthma. *J Appl Physiol* 42: 746–752, 1977.
33. **Persson MG, Wiklund NP, and Gustafsson LE.** Endogenous nitric oxide in single exhalations and the change during exercise. *Am Rev Respir Dis* 148: 1210–1214, 1993.
34. **Phillips CR, Giraud GD, and Holden WE.** Exhaled nitric oxide during exercise: site of release and modulation by ventilation and blood flow. *J Appl Physiol* 80: 1865–1871, 1996.
35. **Pietropaoli AP, Perillo IB, Torres A, Perkins PT, Frasier LM, Utell MJ, Frampton MW, and Hyde RW.** Simultaneous measurement of nitric oxide production by conducting and alveolar airways of humans. *J Appl Physiol* 87: 1532–1542, 1999.
36. **Que LG, Liu L, Yan Y, Whitehead GS, Gavett SH, Schwartz DA, and Stamler JS.** Protection from experimental asthma by an endogenous bronchodilator. *Science* 308: 1618–1621, 2005.
37. **Rouhos A, Ekroos H, Karjalainen J, Sarna S, and Sovijärvi AR.** Exhaled nitric oxide and exercise-induced bronchoconstriction in young male conscripts: association only in atopics. *Allergy* 60: 1493–1498, 2005.
38. **Scollo M, Zanconato S, Ongaro R, Zaramella C, Zacchello F, and Baraldi E.** Exhaled nitric oxide and exercise-induced bronchoconstriction in asthmatic children. *Am J Respir Crit Care Med* 161: 1047–1050, 2000.
39. **Shin HW, Rose-Gottron CM, Cooper DM, Hill M, and George SC.** Impact of high-intensity exercise on nitric oxide exchange in healthy adults. *Med Sci Sports Exerc* 35: 995–1003, 2003.
40. **Shin HW, Rose-Gottron CM, Cooper DM, Newcomb RL, and George SC.** Airway diffusing capacity of nitric oxide and steroid therapy in asthma. *J Appl Physiol* 96: 65–75, 2004.
41. **Shin HW, Rose-Gottron CM, Perez F, Cooper DM, Wilson AF, and George SC.** Flow-independent nitric oxide exchange parameters in healthy adults. *J Appl Physiol* 91: 2173–2181, 2001.
42. **Shin HW, Rose-Gottron CM, Sufi RS, Perez F, Cooper DM, Wilson AF, and George SC.** Flow-independent nitric oxide exchange parameters in cystic fibrosis. *Am J Respir Crit Care Med* 165: 349–357, 2002.
43. **Silkoff PE, McClean PA, Caramori M, Slutsky AS, and Zamel N.** A significant proportion of exhaled nitric oxide arises in large airways in normal subjects. *Respir Physiol* 113: 33–38, 1998.
44. **Silkoff PE, Sylvester JT, Zamel N, and Permutt S.** Airway nitric oxide diffusion in asthma. Role in pulmonary function and bronchial responsiveness. *Am J Respir Crit Care Med* 161: 1218–1228, 2000.
45. **Silkoff PE, Wakita S, Chatkin J, Ansarin K, Gutierrez C, Caramori M, McClean P, Slutsky AS, Zamel N, and Chapman KR.** Exhaled nitric oxide after β_2 -agonist inhalation and spirometry in asthma. *Am J Respir Crit Care Med* 159: 940–944, 1999.
46. **St Croix CM, Wetter TJ, Pegelow DF, Meyer KC, and Dempsey JA.** Assessment of nitric oxide formation during exercise. *Am J Respir Crit Care Med* 159: 1125–1133, 1999.
47. **Storm van's Gravesande K, Wechsler ME, Grasemann H, Silverman ES, Le L, Palmer LJ, and Drazen JM.** Association of a missense mutation in the NOS3 gene with exhaled nitric oxide levels. *Am J Respir Crit Care Med* 168: 228–231, 2003.
48. **Suman OE and Beck KC.** Role of airway endogenous nitric oxide on lung function during and after exercise in mild asthma. *J Appl Physiol* 93: 1932–1938, 2002.
49. **Tee AK and Hui KP.** Effect of spirometric maneuver, nasal clip, and submaximal inspiratory effort on measurement of exhaled nitric oxide levels in asthmatic patients. *Chest* 127: 131–134, 2005.
50. **Terada A, Fujisawa T, Togashi K, Miyazaki T, Katsumata H, Atsuta J, Iguchi K, Kamiya H, and Togari H.** Exhaled nitric oxide decreases during exercise-induced bronchoconstriction in children with asthma. *Am J Respir Crit Care Med* 164: 1879–1884, 2001.
51. **Trolin G, Anden T, and Hedenstierna G.** Nitric oxide (NO) in expired air at rest and during exercise. *Acta Physiol Scand* 151: 159–163, 1994.
52. **Tsoukias NM and George SC.** A two-compartment model of pulmonary nitric oxide exchange dynamics. *J Appl Physiol* 85: 653–666, 1998.
53. **Tsoukias NM, Shin HW, Wilson AF, and George SC.** A single-breath technique with variable flow rate to characterize nitric oxide exchange dynamics in the lungs. *J Appl Physiol* 91: 477–487, 2001.
54. **Venegas JG, Winkler T, Musch G, Vidal Melo MF, Layfield D, Tgavalekos N, Fischman AJ, Callahan RJ, Bellani G, and Harris RS.** Self-organized patchiness in asthma as a prelude to catastrophic shifts. *Nature* 434: 777–782, 2005.
55. **Wechsler ME, Grasemann H, Deykin A, Silverman EK, Yandava CN, Israel E, Wand M, and Drazen JM.** Exhaled nitric oxide in patients with asthma: association with NOS1 genotype. *Am J Respir Crit Care Med* 162: 2043–2047, 2000.
56. **Weibel ER.** *Morphometry of the Human Lung*. New York: Springer, 1963.
57. **Yates DH, Kharitonov SA, Thomas PS, and Barnes PJ.** Endogenous nitric oxide is decreased in asthmatic patients by an inhibitor of inducible nitric oxide synthase. *Am J Respir Crit Care Med* 154: 247–250, 1996.
58. **Yoshikawa T, Shoji S, Fujii T, Kanazawa H, Kudoh S, Hirata K, and Yoshikawa J.** Severity of exercise-induced bronchoconstriction is related to airway eosinophilic inflammation in patients with asthma. *Eur Respir J* 12: 879–884, 1998.